

Management of Pseudomyxoma Peritonei

using eakin **Wound Pouches**[™]

Case study submitted by:
Deb Day, Stomal Therapy CNC Central
Coast Local Health District, Australia

PATIENT HISTORY

- 47-year-old lady;
- Diagnosed with Pseudomyxoma Peritonei 13 years earlier;
- Multiple surgeries and a left-sided, high output ileostomy;
- The lady was very independent and lived at home on total parenteral nutrition (TPN) and intravenous fluids (IVF) via a Hickman's Line;
- Diagnosed disease recurrence of Pseudomyxoma Peritonei in 2014;
- No further surgery;
- Receiving Palliative care.





CARE MANAGEMENT PLAN

- The patient was managed under Palliative Care.
- Continued treatments included:
 - Total Parenteral Nutrition (TPN).
 - Intravenous fluids (IVF).
 - Blood transfusions.
 - Use of a syringe driver for pain relief.
- Pouching the Pseudomyxoma Peritonei with Eakin Wound Pouches™ was deemed appropriate because:
 - They provided comfort, odour control, and containment of wound fluid.
 - They reduced the nurses' workload in providing care due to no additional componentry or reliance on resources such as electricity.



PRODUCT USED

Extra-large eakin **Wound Pouch™**

839265

